



Sepia Ink: An Untouched Molecule from Deep Oceans

Chisti MR¹, Raza A¹, Singh PA¹, Niazi J^{*}

Mohammad Ramjan Chisti, Bahra Institute of Pharmacy, Patiala, India

Aasim Raza, Bahra Institute of Pharmacy, Patiala, India

Preet Amol Singh, Bahra Institute of Pharmacy, Patiala, India

Junaid Niazi, Institute of Pharmacy, Patiala, India

ABSTRACT

Due to ever increasing demand of safer medicine impetus has been on researches in natural products especially marine products as it is still an untouched field. One such product is sepia ink obtained from cuttlefish present in Mediterranean and Atlantic Ocean. It has a very special place in homeopathic medicine and in traditional Chinese medicine along with its culinary uses. It is mainly composed of eumelanin and degradation studies have revealed it is a copolymer of 5, 6-dihydroxyindole and 5,6-dihydroxyindole-2-carboxylic acid. It has tremendous potential in variety of diseases like uterine dysfunction, anticancer, antimicrobial, anti inflammatory activity along with haemopoetic and hepatoprotective effects. Present review aims to assemble various pharmacological activities of sepia ink although deeper insight in its pharmacological potential is required.

Keywords: sepia, cuttle fish, marine drugs, eumelanin

INTRODUCTION

Natural products particularly marine metabolites are increasingly becoming major players in drug discovery due to their structurally exclusive molecules produced by marine organisms. ^{1,2} Recent studies on marine drugs have focused their potential applications in the treatment of human diseases and betterment of human health. Several nautical natural products are currently in pre-clinical and medical evaluation, have shown promising results. ⁽¹⁾ New trends in drug discovery from natural sources emphasize on investigation of the marine ecosystem to explore numerous complex and novel chemical entities. These entities are the sources of new leads for treatment of variety of diseases such as cancer, AIDS, inflammatory conditions including viral, bacterial and fungal infections. ⁽²⁾ Sepia ink mainly composed of melanin has proved to be a valuable substitute and has a wide range of therapeutic applications like anti-radiation, antitumor, immunomodulatory, procoagulant function, etc. ^(3,4,5) Present review aims to include various pharmacological, and phytochemical studies so that better insight on this obscure marine product can be obtained.

SOURCE and Chemistry

Sepia ink is a suspension of melanin granules in a viscous colourless medium obtained from common cuttlefish, *Sepia officinalis* L. (family *Sepiidae*) is distributed in the Mediterranean Sea, Eastern Atlantic-ocean and north-western coast of Africa. ⁽⁶⁾ It is produced at the end of the cuttlefish maturation in digestive tract degenerate. Ejection of dark ink from the sac is a defensive mechanism of cuttlefish employed to avoid dangers and risks. Squid ink is a mixture of large amounts of melanin along with proteins, lipids, glycosaminoglycan's and various minerals. ⁽⁷⁾ Degradation studies on sepia melanin proved that it is a copolymer of 5, 6-dihydroxyindole and 5,6-dihydroxyindole-2-carboxylic acid, which are believed to be main monomeric building blocks of eumelanin. ^(8,9)



TRADITIONAL/ HOMEOPATHIC USES

Nair et al. in a review has ascribed to sepia ink as a traditional medicine, both in Western (ancient Greece and Rome) and Oriental culture (including China).^(10,11) Hahnemann regarded as father of homeopathy reported observations about the cachectic condition of sepia ink which convinced him about its role as antipsoric remedies.⁽¹²⁾ In Chinese traditional medicine it had been reported in the treatment of haemostasis for centuries and listed in the Compendium of Material Medica compiled by Shizhen Li (of Ming Dynasty) for treatment of heart pain and curative effects in gynaecology, surgery, etc.⁽¹¹⁾

PHARMACOLOGICAL ACTIVITIES

Uterine dysfunctioning

In a recent clinical trial conducted by The Institute of Oceanology, China where 400 women with uterine dysfunctional bleeding were given sepia ink capsules has been proved efficient. Improvement has also been seen for the gastrointestinal and chronic tuberculosis bleeding.⁽¹³⁾

Anticancer/Anticytotoxic Effect

The carbohydrate part of peptidoglycan extracted from sepia ink has proved antitumor activity as it linearly decreases DU-145 cell growth in a dose dependent manner although the mechanism of action is still not clear.^(14,15) A study has shown that purified tyrosine with no added extracted from the melanin-free fraction of ink is toxic to transformed human cell lines. Russo et. al., speculated the cytotoxic effect to the production of dopaquinone, which is interacts with nucleophiles to produce protein-bound DOPA through a 5-S-cysteinyl-dopa residue and causes cellular damage by oxidation⁽¹⁶⁾. Peptidoglycans extracted from sepia ink possesses anti-tumour effects as it has been reported that peptidoglycans can affect cell division eukaryotes which include fragmentation of DNA and apoptosis and the reserve of embryonic development.^(17,18,19,20,21) Sulphated sepia ink peptidoglycans derivatives has known to suppress invasion and migration of carcinoma cells by inhibiting matrix metalloproteinase-2.⁽²²⁾ They have also known to suppress melanoma metastasis by inhibiting tumour adhesion and angiogenesis mediated by basic fibroblast growth factor.^(23, 24, 25) Sepia ink oligopeptide has been reported to inhibit prostate cancer cells by inducing apoptosis via activation of caspase-3 and elevation of the ratio of Bax/Bcl-2.^(26, 27, 28)

Hematopoietic Effects

Lie et. al., in 2007 had speculated that cuttlefish ink increases haematopoiesis which results in strengthening of immune responses by it enhancing the proliferation and differentiation of granulocyte-monocyte progenitor cells.⁽²⁹⁾

Anti-Hypertensive Actions

A peptide derivative purified from squid ink causes dilation of blood vessels, resulting in lower blood pressure and acts as a potential anti-hypertensive⁽³⁰⁾.

Anti-Retroviral Activity

Ink from *Loligo duvauceli* and *Sepiella inermis* has been reported to have an anti-retroviral activity.⁽³¹⁾

Anti-ulceration activity

Study conducted in 1982 by Andersen et. al., has concluded that sepia ink possesses marked antiulcer activity by decreasing gastric juice secretion.⁽³²⁾

Anti-Inflammatory Activity

Mimura et al.^(33, 34) reported anti-inflammatory activity for the same fraction of squid ink that inhibits gastric secretion.



Hepatoprotective effects

Studies have shown that sepia ink extract has shown a general decrease in total bilirubin, direct bilirubin and indirect bilirubin in the Bile Duct Ligation (BDL) animal model which is a typical animal model of secondary biliary disease in animals that causes of bile duct epithelial cells proliferation, hepatocellular necrosis, apoptosis, stellate cell activation, and, causes liver fibrosis and cirrhosis and hepatotoxicity.⁽³⁵⁾ Treatment with sepia ink extract caused a decrease in serum gamma glutamyl transferase in rats models due to the protective effect of sepia extract on hepatic injury during bile constriction.⁽³⁶⁾

Antibacterial activity

Ink isolated from *Sepia pharaonis* and purified by column chromatography using diethyl ether showed maximum inhibitory effect against *Pseudomonas aeruginosa*, *Staphylococcus epidermidis*, *Escherichia coli* and *Klebsiella pneumoniae*.⁽³⁷⁾

Haemostatic effect

Chitin and sepia ink hybrid haemostatic sponge has been proven for hemostasis and stimulation of healing. Several kinds of blood parameters were taken into consideration like thrombin time, prothrombin time, activated partial thromboplastin time, fibrinogen and platelet factor 4; anticoagulation parameter including antithrombin III, fibrinolytic parameters including plasminogen, fibrin degradation product etc. Results proved that Chitin and sepia ink hybrid haemostatic sponge has no significant effect on the blood parameters of mice.⁽³⁸⁾

NON PHARMACEUTICAL USE

It is used as food flavouring agent worldwide because of its superior flavour.⁽³⁹⁾ *Arroz negro* (black rice), baby squid in ink sauce, ikasumi jiru ink soup with pork and squid and Caviarne (an imitation caviar) are some of the dishes and foods that use sepia ink. Processed ink is used as a food colouring.⁽⁴¹⁾ Due to its antimicrobial properties cuttlefish ink is also used to cure and, thus, extend the shelf life of cuttlefish meat.^(40,41,42)

CONCLUSION

Sepia ink has acquired unique space in the homeopathy although its not is still not prevalent in the evidence based medicine. Recent researches and studies have shown some of its beneficial pharmacological uses like anticancer, haemostatic, hepatoprotective, antimicrobial activity etc. Detailed work and investigation is required on sepia ink and other marine products as this is the need of hour.

REFERENCES

1. Arya V, Gupta VK. A review on marine immunomodulators. *International journal of pharmacy and life science*, May: 2011, 751-758.
2. Jain R. Marine organisms: Potential source for drug discovery. *Current science*, 2008: 94: 292.
3. Blunt JW, Copp BR, Hu WP, Munro MH, Northcote PT, Prinsep MR. Marine natural products. *Nat Prod Rep*, 2007; 24, 31-86.
4. Bataller R, Brenner DA. Liver fibrosis. *J Clin Invest*, 2005; 115, 209-18.
5. Vate NK, Benjakul S. Antioxidative activity of melanin-free ink from splendid squid (*Loligo formosana*). *International Aquatic Research*, 2013; 5, 9.
6. Khromov DN 1998. Distribution patterns of Sepiidae. In Voss NA, Vecchione M, Toll RB and Sweeney MJ. Systematics and Biogeography of Sepias. *Smith Cont Zool* 586: 191-206.
7. Huazhong Liu, Ping L, Shaohong C, Jianghua S. Effects of Squid Ink on Growth Performance, Antioxidant Functions and Immunity in Growing Broiler Chickens *Asian-Aust. J. Anim. Sci.* 2011; 24 (12), 1752 – 1756.



8. Meredith P, Sarna T. The physical and chemical properties of eumelanin. *Pigment Cell Res.* 2006; 19, 572-594.
9. Guo X, Chen SG, Hu YQ, Li GY, Liao NB, Ye XQ, et. al. Preparation of water-soluble melanin from squid ink using ultrasound assisted degradation and its anti-oxidant activity. *J. Food. Sci. Technol* 2014; 51(12): 3680-3690.
10. Nair JR, Pillai D, Joseph SM, Gomathi P, Senan PV, Sherief PM. Sepia research and bioactive substances. *Indian J. Geo-Mar. Sci.* 2011; 40, 13-27.
11. Zhong JP, Wang G, Shang JH, Pan JQ, Li K, Huang Y, Liu HZ, et. al. Protective effects of squid ink extract towards hemopoietic injuries induced by cyclo-phosphamine. *Marine Drugs* 2009; 7, 9-18.
12. Leiser O, The molluscs: Murex and Sepia, *British Homeopathic Journal* July 1990, Vol. 79, pp. 138-146.
13. Wang XH, Yan YN, Zhang RJA. Comparison of chitosan and collagen sponges as hemostatic dressings *J. Bioact. Compat. Pol.* 2006; 21, 39-54.
14. Takaya Y, Uchisawa H, Hanamatsu K, Narumi F, Okuzuki B, Matsue H. Novel fucose-rich glycosaminoglycans from squid ink bearing repeating unit of trisaccharide structure(-6Gal-NAc α -3GlcA β 1-3Fucal)n. *Biochem Biophysics Res Commun* 1997; 198(2), 560-567.
15. Takaya Y, Uchisawa H, Narumi F, Matsue H. Illexins A, B, and C from squid ink should have a branched structure. *Biochem Biophys Res Commun.* 1996; 226, 335-338.
16. Russo GL, de Nisco E, Fiore G, di Donato P, d'Ischia M, Palumbo A. Toxicity of melanin-free ink of *Sepia officinalis* to transformed cell lines: Identification of the active factor as tyrosinase. *Biochem. Biophys. Res. Commun.* 2003; 308, 293-299.
17. Naraoka T, Chung HS, Uchisawa H, Sasaki J, Matsue H. Tyrosinase activity in antitumor compounds of squid ink. *Food Sci. Technol. Res.* 2000; 6, 171-175.
18. Takaya Y, Uchisawa H, Matsue H, Okuzaki BI, Narumi F, Sasaki JI, Ishida K. An investigation of the antitumor peptidoglycan fraction from squid ink. *Biol. Pharm. Bull.* 1994; 17, 846-849.
19. Sasaki J, Ishita K, Takaya Y, Uchisawa H, Matsue H. Anti-tumour activity of squid ink. *J. Nutr. Sci. Vitaminol.* 1997; 43, 455-461.
20. Kamio M, Grimes TV, Hutchins MH, van Dam R, Derby CD. The purple pigment aplysiocyanin in sea hare ink deters predatory blue crabs through their chemical senses. *Anim. Behav.* 2010; 80, 89-100.
21. Roten CAH, Karamata D. Endogenous synthesis of peptidoglycan in eukaryotic cells. A novel concept involving its essential role in cell division, tumor formation and the biological clock. *Experientia* 1992; 48, 921-931.
22. Priya SV, George MC, Joseph SM, Devika P, Nair JR, Sherief PM. Antiproliferative effect of cuttle fish ink extract on chicken embryo fibroblasts. In Proceedings of the National Seminar on New Frontiers in Marine Bioscience Research National Institute of Ocean Technology, Chennai: Chennai, India, 2004; pp. 35-39.
23. Wang S, Cheng Y, Wang F, Sun L, Liu C, Chen G, et. al. Inhibition activity of sulfated polysaccharide of *Sepiella maindroni* ink on matrix metalloproteinase (MMP)-2. *Biomed. Pharmacother.* 2008; 62, 297-302.
24. Liu C, Li X, Li Y, Feng Y, Zhou S, Wang F. Structural characterisation and antimutagenic activity of a novel polysaccharide isolated from *Sepiella maindroni* ink. *Food Chem.* 2008; 110, 807-813.
25. Zong A, Zhao T, Zhang Y, Song X, Shi Y, Cao H, et al. Anti-metastatic and anti-angiogenic activities of sulfated polysaccharide of *Sepiella maindroni* ink. *Carbohydr. Polym.* 2013; 91, 403-409.



26. Chen S, Xu J, Xue C, Dong P, Sheng W, Yu G, et al. Sequence determination of a non-sulfated glycosaminoglycan-like polysaccharide from melanin-free ink of the squid *Ommastrephes bartrami* by negative-ion electrospray tandem mass spectrometry and NMR spectroscopy. *Glycoconj. J.* 2008; 25, 481–492.
27. Ding GF, Huang FF, Yang ZS, Yu D, Yang YF. Anticancer activity of an oligopeptide isolated from hydrolysates of *Sepia* ink. *Chin. J. Nat. Med.* 2011; 9, 51–55.
28. Huang F, Yang Z, Yu D, Wang J, Li R, Ding G. *Sepia* ink oligopeptide induces apoptosis in prostate cancer cell lines via caspase-3 activation and elevation of Bax/Bcl-2 ratio. *Mar. Drugs* 2012; 10, 2153–2165.
29. Lei M, Wang J, Wang Y, Pang L, Wang Y, Xu W, et. al. Study of the radio-protective effect of cuttlefish ink on hemopoietic injury. *Asia Pac. J. Clin. Nutr.* 2007; 16, 239–243.
30. Kim SY, Kim SH, Song KB. Characterization of an partial purification and angiotensin-converting enzyme inhibitor from squid ink. *Agric. Chem. Biotechnol.* 2003; 46, 122–123.
31. Rajaganapathi J, Thyagarajan SP, Patterson Edward JK. Study on sepia's ink for anti-retroviral activity. *Indian J. Exp. Biol.* 2000; 38, 519–520.
32. Andersen SO, Roepstorff P. Sclerotization of insect cuticle-III. An unsaturated derivative of *N-acetyldopamine* and its role in sclerotization [J]. *Insect Biochem*, 1982; 12(3): 269–276.
33. Mimura T, Itoh S, Tsujikawa K, Nakajima H, Satake M, Kohama Y, et. al. Studies on biological activities of melanin from marine animals. V. Anti-inflammatory activity of low-molecular-weight melanoprotein from squid (Fr. SM II). *Chem. Pharm. Bull.* 1987, 35, 1144–1150.
34. Fahmy SR, Soliman AM. *In vitro* antioxidant, analgesic and cytotoxic activities of *Sepia officinalis* ink and *Coelatura aegyptiaca* extracts. *Afr. J. Pharm. Pharmacol.* 2013; 7, 1512–1522.
35. Cuevas MJ, Tieppo J, Maroni NP, Tuñón MJ, González-Gallego J. Suppression of amphiregulin/epidermal growth factor receptor signals contributes to the protective effects of quercetin in cirrhotic rats. *J Nutr*, 2011; 141, 1299–305.
36. Abu-Bakar A, Arthur DM, Wikman AS, Rahnasto M, Juvonen RO, Vepsäläinen J, et al. Metabolism of bilirubin by human cytochrome P450 2A6. *Toxicology Apply Pharm*, 2012; 261, 50–8.
37. Nithya M, Ambikapathy V, Panneerselvam A. Effect of pharaoh's cuttlefish ink against bacterial pathogens. *Asian Journal of Plant Science and Research*, 2011; 1 (4), 49–55.
38. Zhang W, Yu-Lin S, Dao-Hai C. Effects of chitin and sepia ink hybrid haemostatic sponge on the blood parameters of mice. *Marine Drugs* 2014; 12, 2269–2281.
39. Hazan, M. Essentials of Classic Italian Cooking, Knopf Inc., New York, NY, USA, 1992, pp. 2700–2730.
40. Marquinet A, Inaki J. Process for Producing a Food Colorant, Colorant thus Obtained and Uses Thereof. U.S. Patent 6,329,010 B1, 11 December 2001, 2700–2730.
41. Mochizuki, A. An antiseptic effect of cuttlefish *Sepioteuthis lessoniana* ink. *Nippon. Suisan Gakkaishi* 1979; 45, 1401–1404.
42. Takai M, Yamazaki K, Kawai Y, Inoue N, Shinano H. Effects of squid liver, skin, and ink on chemical characteristics of “ika-shiokara” during ripening process. *Bull. Jap. Soc. Sci. Fish.* 1993; 59, 1609–1615.
43. Xu H, Gou J, Choi GP, Lee HY, Ahn J. Functional properties of squid by-products fermented by probiotic bacteria. *Food Sci. Biotechnol.* 2009; 18, 761–765.